

THE BURAU ESTIMATE FOR THE ENTROPY OF A BRAID

GAVIN BAND AND PHILIP BOYLAND

FEBRUARY 2, 2008

Abstract

The topological entropy of a braid is the infimum of the entropies of all homeomorphisms of the disk which have a finite invariant set represented by the braid. When the isotopy class represented by the braid is pseudo-Anosov or is reducible with a pseudo-Anosov component, this entropy is positive. Fried and Kolev proved that the entropy is bounded below by the logarithm of the spectral radius of the braid's Burau matrix, $B(t)$, after substituting a complex number of modulus 1 in place of t . In this paper we show that for a pseudo-Anosov braid the estimate is sharp for the substitution of a root of unity if and only if it is sharp for $t = -1$. Further, this happens if and only if the invariant foliations of the pseudo-Anosov map have odd order singularities at the strings of the braid and all interior singularities have even order. An analogous theorem for reducible braids is also proved.

1 INTRODUCTION

Artin's braid group and its Burau representation have been extensively studied by many researchers from many points of view. In dynamical applications a braid is often used to describe the motion of a collection of points in the two-dimensional disk. Since the braid depends only on the motion of the points, it is describing an isotopy class of homeomorphisms on the complement of the points. Thus, the interpretation of the braid group on n -strings, B_n , as a mapping class group of the n -punctured disk is frequently used, and so Thurston's classification theorem for surface isotopy classes is an important tool.

The (reduced) Burau matrix, $B(t)$, of a braid $\beta \in B_n$, is an $(n-1) \times (n-1)$ matrix with entries in $\mathbb{Z}[t, t^{-1}]$, i.e. the entries of the matrix are Laurent polynomials over the integers. In the early 1980's two different but closely related dynamical interpretations of the Burau matrix emerged. Using the construction in Franks' paper [Fra81], the Burau matrix can be interpreted as the signed, linking matrix of a certain Axiom A flow associated with the braid. The signed, linking matrix is an enhanced Markov transition matrix which records the linking of the Markov boxes with the strings of the braid as well as the orientations of their images.

The second dynamical interpretation comes from the machinery in Fried's paper [Fri86]. In this case the Burau matrix of a braid arises as the induced action of the associated

mapping class on a $\mathbb{Z}$ -cover, where the first homology of the cover is given the structure of a module over $\mathbb{Z}[t, t^{-1}]$ (this is a standard description, see, for example, [BB05]). In the more general setting of twisted cohomology, Fried showed that after using the appropriate representation of the fundamental group, the spectral radius of the induced action gives a lower bound on the topological entropy. While these interpretations of the Burau matrix were not explicit in either of these two papers, the two authors were certainly aware of them (P.B. personal communication, 1984). See [Boy89] for an introductory exposition of these two interpretations.

Because the topological entropy of a self-map measures a certain kind of exponential growth it is natural to expect that, at least in certain cases, it is detectable from the growth rates of induced maps on various algebraic objects associated with the underlying space. Thus, for example, the growth rate of induced map on first homology (i.e. its spectral radius) gives a lower bound for the topological entropy ([Man74]), as does exponential growth rate of word length in the fundamental group under iteration by f_* ([Bow78, FLP79]).

These lower bounds only depend on the homotopy class of the map, and so it is also natural to ask whether the bounds are attained, i.e. is there a map in the homotopy class that realises the lower bound? For surface homeomorphisms this question was answered by Thurston. One consequence of his classification theorem is that any isotopy class of surface homeomorphisms contains a map with entropy equal to the growth rate on the fundamental group ([Thu88, FLP79]). While this result is invaluable in theory, in practice, the computation of word length growth in non-Abelian groups is very difficult. On the other hand, computations in homology are much more tractable, but frequently give only trivial lower bounds. A fundamental idea in Fried's paper [Fri86] is that there is a middle ground between these two theories provided by the action on twisted homology. In the most concrete case this amounts to providing a systematic way to examine the growth rate of the action on homology in a collection of finite covers.

Thus one sees that the Burau representation provides a lower bound for the topological entropy of the isotopy class represented by a braid. Specifically, if h is a homeomorphism of the n -punctured disk which is represented by the braid $\beta \in B_n$ with Burau matrix $B(t)$, then

$$h_{top}(h) \geq \sup\{\log \text{sr}(B(\eta)) : \eta \in S^1\} \quad (1.1)$$

where $\text{sr}(B(\eta))$ is the spectral radius of the complex matrix $B(\eta)$ obtained by substituting the complex number η with $|\eta| = 1$ into the Burau matrix. This estimate was obtained directly with different methods by Kolev [Kol89]. The estimate in (1.1) and its alternative version in (4.1) below will be called the *Burau estimate*. If the inequality in (1.1) is an equality for $\eta = \eta_0$, then the Burau estimate is said to be *sharp at η_0* .

Since a braid represents an isotopy class, we may use Thurston's classification scheme to classify braids. Thus a braid is said to be pseudo-Anosov, finite order or reducible if its corresponding isotopy class is. If a braid is finite order or reducible with all finite order components, there is a map in the class with zero topological entropy, and so the Burau estimate is already sharp at $\eta = 1$. The main result here for pseudo-Anosov braids is

Theorem 1.1 *For a pseudo-Anosov braid β , the Burau estimate is sharp at some root of*

unity η_0 if and only if it is sharp at -1 . Further, this happens if and only if the invariant foliations for a pseudo-Anosov-map in the class represented by β have odd order singularities at all punctures and all interior singularities are even order.

A portion of this theorem was obtained in [SKL02] (see Remark 5.2 below). The substitution of complex numbers on the unit circle which are not roots of unity requires different methods. In a subsequent paper we will show that for a pseudo-Anosov-braid, $\text{sr}(B(e^{2\pi i\theta})) < \lambda$ for all $\theta \notin \mathbb{Q}$.

There is an analogous theorem for reducible braids with at least one pseudo-Anosov component. Its statement is rather complicated (see Theorem 6.2 below), but one useful consequence is the following.

Theorem 1.2 *For any braid β on n strings, the Burau estimate is sharp at some root of unity η_0 if and only if it is sharp at a k -th root of unity $e^{2\pi ij/k}$, for some $k \leq \frac{2}{3}n$ which is a power of 2.*

There are two main components in the proof of these theorems. The main algebraic tool is Lemma 3.2 which shows that the union of the spectra obtained by substituting all the k^{th} roots of unity into the Burau matrix $B(t)$ yields essentially the entire spectrum of the action on first homology of a lift of the corresponding mapping class to the k -fold cover. This is coupled with information about the connection of entropy, the Thurston normal form, the action on homology, and the orientability of the invariant foliations of a pseudo-Anosov-map.

The investigations of this paper were partly inspired by recent work using the Burau estimate to get lower bounds on the entropy of fluid flows by applying the Burau estimate to the braids generated by large collections of points moving with the fluid ([Thi05, GFT06, TF06]). For these applications a good understanding of “sharpness” of the estimate is necessary and this paper provides a first step. In addition, questions surrounding the Burau representation provide an important, special case of the more general question of dynamics on abelian covers of surfaces which we investigate in a subsequent paper.

2 PRELIMINARIES

2.1 STANDING HYPOTHESES AND CONVENTIONS

In this paper surfaces X will always be orientable, perhaps with boundary, and compact except perhaps for a finite number of punctures. We fix a Riemannian metric on X which allows us to speak of the lengths of tangent vectors. Self-homeomorphisms of the surface $f : X \rightarrow X$ are always orientation-preserving. If no coefficient ring for homology is specified, it is assumed to be the integers $\mathbb{Z}$, and so $H_1(X) := H_1(X; \mathbb{Z})$. The induced map of the homeomorphism f on first homology is denoted f_* . If M is a square, complex matrix, then $\text{sr}(M)$ denotes its spectral radius.

The classification theorem for regular connected covering spaces identifies each such cover with a normal subgroup of the fundamental group of the connected base space X , or equivalently, with an epimorphism $\pi_1(X) \rightarrow G$, where G is the group of deck transformations of the cover (see, for example, [Ful95]). In this paper G will always be abelian, and so we often designate a cover $\tilde{X}$ by an epimorphism $\rho : H_1(X) \rightarrow G$, with the Hurewicz homomorphism

$\pi_1(X) \rightarrow H_1(X)$ being implicit.

More generally, we shall also need to consider disconnected covers over connected and disconnected base spaces. In these cases it will be convenient to continue to designate the cover by a homomorphism $\rho : H_1(X) \rightarrow G$, which perhaps is not surjective. As this is less commonly encountered, we describe the cover $\tilde{X}$ associated to such a homomorphism ρ . First suppose X is connected, and let $G' = \text{im } \rho \subset G$. Then ρ induces an epimorphism $\rho' : H_1(X) \rightarrow G'$ with the same domain as ρ but with range G' , and so as above it determines a connected covering space $\tilde{X}_0$ over X with deck group G' . We define $\tilde{X}$ to be the disjoint union of copies of $\tilde{X}_0$, one such copy for each coset of G' in G . The action of G' by deck transformations on each copy of $\tilde{X}_0$ extends to an action of G on $\tilde{X}$, in such a way that for any $g, g' \in G$, the element g sends the copy of $\tilde{X}_0$ corresponding to $g' + G$ to that corresponding to $g + g' + G$. Then $\tilde{X}/G \cong X$, and so this makes $\tilde{X}$ a covering space over X with deck group G . Finally, if X is disconnected, we define $\tilde{X}$ to be the disjoint union, over all connected components Y of X , of the covering space $\tilde{Y}$ corresponding to $\rho|_{H_1(Y)}$ as above.

In all cases every homeomorphism of the base $f : X \rightarrow X$ will satisfy $\rho f_* = \rho$, which implies that f lifts to $\tilde{f} : \tilde{X} \rightarrow \tilde{X}$ which commutes with all deck transformations $g \in G$.

2.2 THE BRAID GROUP AND THE BURAU REPRESENTATION

In this section we briefly survey relevant results from the point of view of Dynamical Systems. The classic reference are [Art25, Bir74], and see [BB05] for a survey including recent developments and [Boy94] for a survey of dynamical applications. The book [Fen83] and the classic paper [Mil68] are good sources of information on homology of cyclic covers.

The braid group on n strings, B_n , is defined using generators and relations as

$$B_n = \langle \sigma_1, \dots, \sigma_{n-1} \mid \sigma_i \sigma_k = \sigma_k \sigma_i \text{ if } |i - k| > 2, \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} \rangle.$$

In this paper we shall be principally concerned with B_n interpreted as a mapping class group, namely, the group of isotopy classes of homeomorphisms of the n -punctured disk where all homeomorphisms and all isotopies are required to fix the boundary pointwise. Letting $x_j = j/(n+1)$ for $j = 1, \dots, n$ and $D_n = \{z \in \mathbb{C} : |z - 1/2| \leq 1/2\} \setminus \{x_1, \dots, x_n\}$, the generator σ_i of B_n corresponds to a homeomorphism that switches x_i and x_{i+1} in a counter-clockwise direction. When we indicate a braid $\beta \in B_n$ we will always be identifying it with its corresponding isotopy class, and so for example, $h \in \beta \in B_n$ means that h is a homeomorphism of D_n that is contained in the isotopy class corresponding to β .

As with the braid group we shall need an interpretation of the Burau representation with dynamical content, as the action on homology in a particular cover. To construct the $\mathbb{Z}$ -cover relevant to the Burau representation, we fix a basepoint $x_0 \in D_n$, and around each puncture x_i we take a small clockwise loop Γ_i which we then homotope so it begins and ends at x_0 . Then $\pi_1(D_n, x_0)$ and $H_1(D_n)$ are freely generated by the set $\{\gamma_i\}$ of homotopy/homology classes of the Γ_i 's. Let τ be the epimorphism of $H_1(D_n)$ onto $\mathbb{Z}$ generated by $\tau(\gamma_i) = 1$ for all i ; the resulting cover is called the *Burau cover* and is denoted $D^{(\infty)}$. By construction the deck group of $D^{(\infty)}$ is isomorphic to $\mathbb{Z}$, and we call its generator T . An orientation-preserving homeomorphism h of D_n must permute the punctures of D_n and will therefore act as a permutation on the generators of $H_1(D_n)$, and so $\tau h_* = \tau$. Thus any orientation-preserving

homeomorphism h of D_n will lift to a homeomorphism $\tilde{h} : D^{(\infty)} \rightarrow D^{(\infty)}$ and further, each lift of h commutes with all deck transformations, or $T^j \tilde{h} = \tilde{h} T^j$ for all $j \in \mathbb{Z}$. Note that by definition, any $h \in \beta \in B_n$ is a homeomorphism that fixes the outside boundary of D_n point-wise and this yields a preferred lift of h to $D^{(\infty)}$, namely, the lift which fixes the lift of the outside boundary of D_n point-wise. Unless indicated otherwise any lift to $D^{(\infty)}$ of an $h \in \beta$ will be this preferred lift.

Next note that the first homology group $H_1(D^{(\infty)})$ has a natural structure as a module over $R := \mathbb{Z}[t^{\pm 1}]$, the ring of all Laurent polynomials with coefficients in $\mathbb{Z}$, i.e. the group ring of $\mathbb{Z}$. To describe this structure, lift all the loops Γ_i to arcs $\tilde{\Gamma}_i \in D^{(\infty)}$ with all $\tilde{\Gamma}_i$ starting at some point $\tilde{x}_0$ and ending at the point $T\tilde{x}_0$. Thus for $i = 1, \dots, n-1$, $\zeta_i := [\tilde{\Gamma}_{i+1} - \tilde{\Gamma}_i] \in H_1(D^{(\infty)})$. For a Laurent polynomial $p(t) = \sum a_j t^j \in R$, $p(t)\zeta_i$ represents the class $\sum a_j T^j \zeta_i \in H_1(D^{(\infty)})$, and so as an R -module, $H_1(D^{(\infty)}) \cong R^{n-1}$. Now since the lift of a homeomorphism, $\tilde{h}$, commutes with T , we see that that $\tilde{h}$ acts on $H_1(D^{(\infty)})$ by an R -module isomorphism. So with respect to the R -module basis $\{\zeta_i\}$ of $H_1(D^{(\infty)})$, $\tilde{h}$ acts by a matrix $B(\tilde{h}) \in \text{GL}(n-1, R)$.

For a braid $\beta \in B_n$, pick $h \in \beta$ and its preferred lift $\tilde{h}$ to $D^{(\infty)}$. The matrix $B(\beta) = B(\tilde{h})$ is called the *reduced Burau matrix* of β , and the corresponding homomorphism $B_n \rightarrow \text{GL}(n-1, R)$ the *reduced Burau representation* of the braid group B_n . When the braid β is fixed, we often will write its Burau matrix as $B(t)$. The full Burau representation will not be used here, but for completeness we note that it can be defined similarly using the action of $\tilde{h}$ on the relative homology group $H_1(D^{(\infty)}, F)$, where F is the fiber above the basepoint x_0 (which in this case must be a fixed point of h).

Since our results work for all $n > 2$, we fix once and for all such an n and suppress the dependence of objects on n when possible. So, for example, we write just D not D_n .

2.3 THE NIELSEN-THURSTON NORMAL FORM

Since we have identified the braid group B_n with a surface mapping class group, Thurston's classification theorem will be of central importance here. This theorem identifies "simplest" representatives in any isotopy class. See [FLP79] and [Thu88] for more information. There are minor differences in the literature in how punctures, boundary and the reducible case are handled in stating Thurston's results. The version we give in Theorem 2.1 is adapted to our use with the braid group.

The two main ingredients in Thurston's classification are finite order and pseudo-Anosov maps. A map ϕ is *finite order* if $\phi^n = \text{id}$ for some $n \geq 1$. The map ϕ is pseudo-Anosov if there are a pair of transverse, ϕ -invariant measured foliations¹, $\mathcal{F}^u$ and $\mathcal{F}^s$. Under the action of ϕ the transverse measures are expanded and contracted by a number $\lambda > 1$, which is called the *expansion factor* of the pseudo-Anosov map. This fact is usually indicated by the notation $\phi_* \mathcal{F}^u = \lambda \mathcal{F}^u$ and $\phi_*^{-1} \mathcal{F}^s = \lambda \mathcal{F}^s$. The supporting surface of a pseudo-Anosov map ϕ may be connected or disconnected, but in the latter case we require that ϕ cyclically permutes the components.

Since the structure of the singular points in a measured foliation is central here we

¹The designation "measured foliation" is the standard shortening of the more proper, and much longer name, "foliation with conical singularities with a holonomy invariant transverse measure".




Figure 1: (a) Constructing prongs; (b) a boundary three prong.

describe it in more detail. Near a *regular point* a measured foliation looks locally like the foliation of $\mathbb{R}^2$ by horizontal lines. A measured foliation is also allowed to have a finite number of non-regular or singular points which are required to have a very specific local structure which is characterized by the order of the singularity, i.e. by the number of leaves coming directly into the point. The local structure can be succinctly described using covers branched over $0 \in \mathbb{C}$. Starting with the foliation of $\mathbb{C}$ by horizontal lines and projecting under $z \mapsto z^2$, we get the local structure of an *order one or one-prong singularity* at 0 (see Figure 1(a)). Lifting the one-prong by the map $z \mapsto z^n$ for $n > 2$ gives an *order n or n -prong singularity*. An *order n -punctured singularity* is formed by removing the singular point from an n -prong, and an *order n -boundary singularity* is obtained by replacing this puncture with a boundary circle (see Figure 1(b)).

The invariant foliations associated with a pseudo-Anosov map have a few special additional qualifications. They can have punctured or boundary one-prongs, but interior one-prongs do not occur as they will not persist under isotopy. An interior two-prong is a regular point and is not considered a singularity, but a punctured or boundary two-prong is considered a singularity. Given a measured foliation $\mathcal{F}$ on a surface X with genus g , the Euler-Poincaré-Hopf formula says that

$$2 - 2g = \sum (1 - \kappa_i/2), \quad (2.1)$$

where the sum is over singularities of all types (interior, boundary and punctured) of $\mathcal{F}$, and κ_i is the order of the i^{th} singularity.

Now recall that any homeomorphism $h \in \beta \in B_n$ must fix the boundary of D_n pointwise, a property that is not shared by the Thurston representatives under the usual definitions. In addition, the isotopies used in the braid group must fix the boundary pointwise. These two facts require a small alteration in the designation of Thurston representative in an isotopy class. Here is a version of Thurston's classification theorem adapted to our situation. It follows, for example, by altering the constructions in [Boy99].

Theorem 2.1 (Thurston) *Let f be a homeomorphism of the possibly disconnected, possibly bordered surface X , compact except for a finite number of punctures, and assume that f fixes the boundary of X pointwise. Then there is a homeomorphism Φ isotopic to f by an isotopy*

which fixes the boundary pointwise, and a decomposition

$$X = A \cup \bigcup_{j=1}^m X_j$$

of X into pairwise disjoint Φ -invariant sets, with the following properties.

1. A is a finite disjoint union of embedded open annuli. The waist curve of such an annulus is never null-homotopic; nor are the waist curves of two annuli mutually homotopic. If the waist curve of an annulus $a \in A$ is homotopic to a boundary component b of X , then b is a component of ∂a (and by convention we include b in a).
2. Each X_j is the union of a collection of connected components of $X \setminus A$ which are permuted cyclically by Φ ; and the restriction of Φ to X_j is either finite order or pseudo-Anosov.
3. The restriction of Φ to A has zero topological entropy.

2.4 BRANCHED COVERING SPACES AND ORIENTED FOLIATIONS

Given a finite set $B \subset X$, the triple $p : Y \rightarrow X$ is called a *covering space branched over B* if p is onto and restricts to a covering map (i.e. a surjective local homeomorphism which is evenly covered over any small neighborhood, c.f. [Ful95]) of $Y \setminus p^{-1}B$ over $X \setminus B$. Now let $\mathcal{F}$ be a measured foliation on the surface X . The foliation $\mathcal{F}$ is *orientable* provided there is a vector field Υ on X , zero at the singularities of $\mathcal{F}$, and everywhere else nonzero and tangent to $\mathcal{F}$. When $\mathcal{F}$ is not orientable, it is often useful to orient it by lifting it to a two-sheeted branched cover constructed as follows. Start by puncturing X at the singularities of the foliation and denote the resulting space by X' . Now define $\vec{X}'$ as the space of unit tangent vectors to $\mathcal{F}$ in X' with the topology induced as a subspace of the unit tangent bundle of X . It is evident that $\vec{X}'$ is a two-sheeted cover over X' . By sewing back in the punctures, we obtain a two-fold branched cover $\vec{p} : \vec{X} \rightarrow X$, and pulling back $\mathcal{F}$, we obtain a measured foliation $\vec{\mathcal{F}}$ on $\vec{X}$. The space $\vec{X}$ equipped with the pulled back foliation $\vec{\mathcal{F}}$ is called the *orientation double-cover* of the non-orientable foliation $\mathcal{F}$.

If γ is a smooth loop in X' , then since X is an orientable surface, the tangent bundle restricted to γ is trivial and so there is a well defined monodromy as we pull along γ the unit vectors tangent to $\mathcal{F}$. If this monodromy brings a vector back to itself, we say that *the foliation is oriented along γ* , and if it brings a vector back to its opposite, *the foliation is disoriented along γ* . It is evident from the definition of the orientation double cover that in the first case γ lifts to a pair of disjoint loops in $\vec{X}$, while in the latter case, γ is covered by a single loop $\tilde{\gamma} \subset \vec{X}$, and the covering map induces a degree-two map $\tilde{\gamma} \rightarrow \gamma$. Thus a foliation is oriented if and only if it is oriented along every loop in the complement of the singularity set, and if $\vec{p} : H_1(X') \rightarrow \mathbb{Z}_2$ is the epimorphism associated to the cover $\vec{X}$, then the foliation is oriented along γ if and only if $\vec{p}([\gamma]) = 0$.

The next lemma gives a simple criterion for checking when the foliation is oriented when lifted to a cover. Its proof is standard and we omit it. Implicit in the statement of (ii) is

the fact that if γ is a small loop surrounding a singularity P , then $\vec{\rho}([\gamma]) = 0$, if P has even order, and $\vec{\rho}([\gamma]) = 1$, if P has odd order. Thus if all interior singularities of a measured foliation on X are of even order, then we may treat $\vec{\rho}$ as being defined on $H_1(X)$.

Lemma 2.2 *Assume that X is a possibly disconnected, possibly bordered surface X , compact except for finitely many punctures, and that $\mathcal{F}$ is a measured foliation on X . Let $\rho : H_1(X) \rightarrow G$ be a homomorphism of $H_1(X)$ to a finite abelian group G and let $\tilde{X}$ be the corresponding covering space of X . The following are equivalent:*

- (i) *The lift of $\mathcal{F}$ to $\tilde{X}$ is orientable;*
- (ii) *All singularities in the interior of X have even order, and there exists a homomorphism $\delta : \text{im } \rho \rightarrow \mathbb{Z}_2$ such that $\delta \circ \rho = \vec{\rho}$, where $\vec{\rho}$ is the morphism defining the orientation cover of $\mathcal{F}$.*

3 FINITE COVERS AND SUBSTITUTING ROOTS OF UNITY

In this section we show that the substitutions of k^{th} roots of unity into the Burau representation of a braid give all the essential spectral information about the action of the braid's mapping class on homology in the k -fold cover. The reducible case considered in Section 6 requires us to work with more general subsurfaces of D and, in fact, the results of this section apply to fairly general topological spaces.

Let $h : X \rightarrow X$ be a homeomorphism of the perhaps disconnected space X . Suppose that $\rho : H_1(X) \rightarrow \mathbb{Z}$ is some homomorphism which satisfies $\rho h_* = \rho$, and denote by $p^{(\infty)} : X^{(\infty)} \rightarrow X$ the covering space associated to ρ as in Section 2.1. The covering $p^{(\infty)}$ is thus generated by a deck group isomorphic to $\mathbb{Z}$, and we denote the generator of this deck group by T . Let $h^{(\infty)}$ be a lift of h to $X^{(\infty)}$. The condition $\rho h_* = \rho$ implies that $h^{(\infty)}$ commutes with T and hence also with every other element of the deck group.

The main example for our purposes is where $X = D$ is the n -punctured disk, $h : D \rightarrow D$ is a representative of the braid $\beta \in B_n$, and $\rho = \tau$ is the homomorphism defining the Burau cover $D^{(\infty)}$ of D (see Section 2.2). To deal with reducible braids we will also have to consider certain h -invariant subsurfaces of D (which may be disconnected). In general, all we shall assume about the space X is that its first homology group $H_1(X)$ is free and of finite rank. The results can easily be extended to work whenever the torsion-free part of $H_1(X)$ has finite rank.

To construct cyclic covers, for each integer $k > 0$, let $\xi_k : \mathbb{Z} \rightarrow \mathbb{Z}_k$ be the quotient homomorphism and define $\rho_k = \xi_k \circ \rho$. Let $p^{(k)} : X^{(k)} \rightarrow X$ be the covering space associated to ρ_k , so $X^{(k)} = X^{(\infty)}/T^k$. Let $q^{(k)} : X^{(\infty)} \rightarrow X^{(k)}$ be the covering projection. The image under $q_*^{(k)}$ of $H_1(X^{(\infty)})$ is a subgroup of $H_1(X^{(k)})$ which we denote by $S^{(k)}$.

The map $h^{(\infty)}$ pushes down to a well-defined lift $h^{(k)}$ of h on $X^{(k)}$. Also, the deck group generator T of $X^{(\infty)}$ pushes down to a generator of the deck group for $p^{(k)}$, which we also denote by T . Both $h_*^{(k)}$ and T_* leave $S^{(k)}$ invariant by definition.

As in Section 2.2, we write $R = \mathbb{Z}[t^{\pm 1}]$ for the ring of all Laurent polynomials with coefficients in $\mathbb{Z}$ (i.e. the group ring of $\mathbb{Z}$) and we treat the integral homology group $H_1(X^{(\infty)})$ as a module over R . The next lemma generalizes Section 2.2 to show that, just as for the

case of the Burau cover $D^{(\infty)}$, the first homology of $X^{(\infty)}$ is a free module of finite rank over R .

Lemma 3.1 *Suppose $H_1(X)$ is a free Abelian group of finite rank r_1 , and let r_0 be the number of connected components $X_0 \subseteq X$ such that $\rho|_{H_1(X_0)}$ is not identically 0. Then $H_1(X^{(\infty)})$ is a free R -module of rank $r = r_1 - r_0$.*

Proof: Let $X' \subset X$ be a set consisting of one point in each connected component of X , and let $Z = (p^{(\infty)})^{-1}(X')$. Let $\gamma_1, \dots, \gamma_{r_1}$ be a collection of loops in X , each based at some point of X' , whose homology classes generate $H_1(X)$, and lift each γ_j to a path $\tilde{\gamma}_j$ in $X^{(\infty)}$. Because $X^{(\infty)}$ is a covering space with deck group $\{T^i\}$, the homology classes of paths of the form $T^i \tilde{\gamma}_j$ generate the relative homology group $H_1(X^{(\infty)}, Z)$ as a vector space. In other words, the homology classes of the $\tilde{\gamma}_j$'s generate $H_1(X^{(\infty)}, Z)$ as an R -module. Since $H_1(X)$ is free, it follows that there can be no torsion in this R -module, and therefore $H_1(X^{(\infty)}, Z)$ is free and of rank r_1 . Moreover, since Z is discrete, the inclusion $(X^{(\infty)}, \emptyset) \subset (X^{(\infty)}, Z)$ induces an injection $H_1(X^{(\infty)}) \rightarrow H_1(X^{(\infty)}, Z)$ of R -modules, so $H_1(X^{(\infty)})$ is a free R -module of rank at most r_1 . The formula $r = r_1 - r_0$ can now be proved by considering the long exact homology sequence of the pair $(X^{(\infty)}, Z)$. ■

Let $\zeta_1, \dots, \zeta_r$ be a basis for $H_1(X^{(\infty)})$. With respect to this basis, the module isomorphism $h_*^{(\infty)}$ is given by a square matrix $M = M(t) \in \text{GL}(r, R)$ with entries in R . Note that M depends on the morphism ρ used to define $X^{(\infty)}$, on the homotopy class of h , and also on the choice of lift of h to $X^{(\infty)}$. If $X = D$ is the n -punctured disk, h is a representative the braid β on n strings, ρ is the homomorphism τ of Section 2.2 and $h^{(\infty)}$ is the preferred lift of h to $D^{(\infty)}$, then we have $M = B(\beta)$, the reduced Burau matrix of β .

For a complex number $\eta \in \mathbb{C}$, denote by $M(\eta)$ the complex matrix obtained from M by substituting η in place of t . If $\eta \neq 0$ then this matrix is invertible, just as M is, and so it acts as a linear isomorphism of $\mathbb{C}^r$ to itself. We denote by $S_{\mathbb{C}}^{(k)}$ the complexification of $S^{(k)}$ and treat $S_{\mathbb{C}}^{(k)}$ as a subspace of $H_1(X^{(k)}, \mathbb{C})$.

The next lemma connects the action on the k -fold cover to substitutions of complex k^{th} roots of unity into the matrix $M(t)$. It is based on well-known, elementary facts. Depending on the chosen perspective, it follows, for example, from the splitting of a representation of Z_k into the sum of irreducibles, or from the invertibility of the order- k discrete Fourier transform. Rather than abstract the necessary algebra and then apply it to the situation at hand, it is simpler to maintain an algebraic topology perspective and give a direct proof using the invariance of an eigen-decomposition.

Lemma 3.2 *Let T be the generator of the deck group for the covering $p^{(k)} : X^{(k)} \rightarrow X$, and let $h^{(k)}$ and $h^{(\infty)}$ be the lifts of h to $X^{(k)}$ and $X^{(\infty)}$. The eigenvalues of T_* restricted to $S_{\mathbb{C}}^{(k)}$ are $1, \eta_k, \eta_k^2, \dots, \eta_k^{k-1}$ where $\eta_k = e^{2\pi i/k}$. Denote by $E_0, \dots, E_{k-1}$ the corresponding eigenspaces in $S_{\mathbb{C}}^{(k)}$. Then each subspace E_m is $h_*^{(k)}$ -invariant, and the action of $h_*^{(k)}$ on E_m is given by the matrix $M(\eta_k^m)$, obtained by substituting η_k^m into the matrix $M(t)$ of $h_*^{(\infty)}$.*

Proof: Let $R^{(k)}$ be the ring of all Laurent polynomials in a variable s which satisfies $s^k = 1$, so $R^{(k)}$ is isomorphic to the group ring $\mathbb{Z}[Z_k]$. As with $H_1(X^{(\infty)})$, we can treat $H_1(X^{(k)})$

as a free module over $R^{(k)}$. Note that T_* and $h_*^{(k)}$ act by module isomorphisms, and $S^{(k)}$ is an $R^{(k)}$ -submodule that is invariant under both T_* and $h_*^{(k)}$.

Since T_* has order k its eigenvalues are as given. A general element of $S_{\mathbb{C}}^{(k)}$ has the form $\sum_{i \in \mathbb{Z}_k} s^i \left(\sum_{j=1}^{n-1} a_{i,j} \zeta_j^{(k)} \right)$, for complex numbers $a_{i,j} \in \mathbb{C}$. For $m = 0, \dots, k-1$, let E_m be the set of elements which when written in this form satisfy $a_{i+1,j} = \eta_k^{-m} a_{i,j}$ for each $i \in \mathbb{Z}_k$ and all j . Since T_* acts as an $R_{\mathbb{C}}^{(k)}$ -module homomorphism on $S_{\mathbb{C}}^{(k)}$ by multiplication by s , the set E_m consists of eigenvectors of T_* with eigenvalue η_k^m . Further, the dimension of E_m as a complex vector space is r and so $E_0 \oplus \dots \oplus E_{k-1} = S_{\mathbb{C}}^{(k)}$.

Since $h_*^{(k)}$ commutes with T_* , each E_m is $h_*^{(k)}$ -invariant. By definition, the matrix $M \in \text{GL}(r, R)$ is the matrix of $h_*^{(\infty)} : H_1(X^{(\infty)}) \rightarrow H_1(X^{(\infty)})$ relative to the chosen basis $\{\zeta_i\}$ of $H_1(X^{(\infty)})$. We decompose this matrix as

$$M = \sum_{i \in \mathbb{Z}} t^i M_i,$$

with each $M_i \in \text{Mat}(r, \mathbb{Z})$. Projecting this action to $S_{\mathbb{C}}^{(k)}$ we have that $h_*^{(k)}$ acts as an $R_{\mathbb{C}}^{(k)}$ -module homomorphism on $S_{\mathbb{C}}^{(k)}$ by the matrix

$$M(s) := \sum_{i \in \mathbb{Z}_k} s^i \left(\sum_{l \in \mathbb{Z}} M_{lk+i} \right) \in \text{GL}(r, R^{(k)}). \quad (3.1)$$

On the other hand, since $\eta_k^k = 1$, the matrix $M(\eta_k^m)$ is given by

$$M(\eta_k^m) = \sum_{i \in \mathbb{Z}} \eta_k^{mi} M_i = \sum_{i=0}^{k-1} \eta_k^{mi} \left(\sum_{l \in \mathbb{Z}} M_{lk+i} \right). \quad (3.2)$$

If $v \in E_m$ then by (3.1) and (3.2) we have

$$\begin{aligned} h_*^{(k)}(v) &= \sum_{i \in \mathbb{Z}_k} s^i \left(\sum_{l \in \mathbb{Z}} M_{lk+i} \right) \cdot v \\ &= \sum_{i \in \mathbb{Z}_k} \left(\sum_{l \in \mathbb{Z}} M_{lk+i} \right) \cdot \eta_k^{mi} v \\ &= M(\eta_k^m) \cdot v \end{aligned}$$

as claimed. ■

Although $H_1(X^{(k)})$ is larger than $S^{(k)}$, the next lemma indicates that all of the *growth* of $h_*^{(k)}$ on $H_1(X^{(k)})$ occurs in $S^{(k)}$.

Lemma 3.3 *With notation as above, the eigenvalues of $h_*^{(k)}$ acting on $H_1(X^{(k)})$ are those of its restriction to $S^{(k)}$ together with some roots of unity. In particular, the spectral radius of $h_*^{(k)}$ equals that of its restriction to $S^{(k)}$.*

Proof: We return to treating $H_1(X^{(k)})$ and $S^{(k)}$ as Abelian groups. Since $S^{(k)}$ is $h_*^{(k)}$ -invariant, we may write $h_*^{(k)}$ as a matrix of the form

$$h_*^{(k)} = \begin{pmatrix} A & B \\ 0 & C \end{pmatrix}$$

in which A represents the action of $h_*^{(k)}$ on $S^{(k)}$, and C its action on $h_*^{(k)}$ on $H_1(X^{(k)})/S^{(k)}$. We will construct a basis for $H_1(X^{(k)})/S^{(k)}$ whose elements are permuted by $h_*^{(k)}$. In particular the eigenvalues of C are all roots of unity, so this will prove the lemma.

To construct the desired basis, let Y be a connected component of $X^{(k)}$ and let $X_0 = p^{(k)}(Y)$. The image of $H_1(X_0)$ under the morphism $\rho : H_1(X) \rightarrow \mathbb{Z}$ defining $X^{(\infty)}$ is some subgroup $a\mathbb{Z}$ of $\mathbb{Z}$. The chosen connected component Y then corresponds to a coset of $a\mathbb{Z}_k$ in $\mathbb{Z}_k$, and so the deck transformations which preserve Y are precisely those of the form T^i with $i \in a\mathbb{Z}_k$.

Choose a loop γ in X_0 such that $\rho([\gamma]) = a$. This loop γ may lift in more than one way to a path in Y . Let $\tilde{\gamma}$ be one such lift. We define an element $\gamma(Y) \in H_1(X^{(k)})/S^{(k)}$ by

$$\gamma(Y) = \left[\sum_{i \in a\mathbb{Z}_k} T^i \tilde{\gamma} \right] + S^{(k)}.$$

where, as usual, the square brackets denote taking the homology class. Any other lift of γ to Y is of the form $T^i \tilde{\gamma}$ for $i \in a\mathbb{Z}_k$, so $\gamma(Y)$ does not depend on the choice of lift of γ .

We claim that $\gamma(Y)$ also does not depend on the chosen loop γ . For suppose that γ' is another loop in X_0 satisfying $\rho([\gamma']) = a$. Then $\rho(\gamma - \gamma') = 0 \in \mathbb{Z}$, so the closed 1-chain $\gamma' - \gamma$ lifts to a closed 1-chain in $X^{(\infty)}$, representing an element of $H_1(X^{(\infty)})$. If the lift $\tilde{\gamma}'$ of γ' is chosen so that $\tilde{\gamma}' - \tilde{\gamma}$ is also a closed 1-chain in $X^{(k)}$, then it follows that $[\tilde{\gamma}' - \tilde{\gamma}] \in S^{(k)}$. Therefore also $\sum_i T^i(\tilde{\gamma}' - \tilde{\gamma}) \in S^{(k)}$, proving that $\gamma'(Y) - \gamma(Y) = 0 \in H_1(X^{(k)})/S^{(k)}$.

Note that if $a = 0$ then $\gamma(Y) = 0 \in H_1(X^{(k)})/S^{(k)}$, since in that case Y lifts to a component of $X_0^{(\infty)}$ homeomorphic to X_0 . On the other hand if $a \neq 0$ then the deck group of any connected component of $X_0^{(\infty)}$ over X_0 is infinite cyclic, so no lift of $\sum_{i \in a\mathbb{Z}_k} T^i \tilde{\gamma}$ can represent a homology class in $H_1(X^{(\infty)})$. Hence $\gamma(Y) = 0$ iff $a = 0$. A dimension calculation similar to that in Lemma 3.1 now shows that the set

$$\{\gamma(Y) | Y \text{ is a connected component of } X^{(k)}\}$$

generates $H_1(X^{(k)})/S^{(k)}$. To finish the proof, it is enough to note that, since $h^{(k)}$ permutes the connected components of $X^{(k)}$, and since $\rho h_* = \rho$, the map $h_*^{(k)}$ must also permute the $\gamma(Y)$'s. ■

Lemma 3.3 together with Lemma 3.2 yields

Theorem 3.4 *Let $h : X \rightarrow X$ be a homeomorphism of the locally path-connected, semi-locally simply connected topological space X , whose first homology group we assume to be free and of finite rank. Suppose $\rho : H_1(X) \rightarrow \mathbb{Z}$ is a homomorphism which satisfies $\rho h_* = \rho$, and let $X^{(\infty)}$ and $X^{(k)} = X^{(\infty)}/T^k$ denote the covering spaces over X corresponding to ρ*

and $\xi_k \circ \rho$, with covering projection $q^{(k)} : X^{(\infty)} \rightarrow X^{(k)}$. Let $h^{(\infty)}$ and $h^{(k)}$ denote lifts of h to these covering spaces. If $M = M(t) \in GL(r, R)$ denotes the matrix of $h_*^{(\infty)} : H_1(X^{(\infty)}) \rightarrow H_1(X^{(\infty)})$ as an R -module isomorphism, then the action of $h_*^{(k)}$ on the invariant subspace $S_{\mathbb{C}}^{(k)} = q_*^{(k)}(H_1(X^{(\infty)}, \mathbb{C}))$ is given by the direct sum

$$h_*^{(k)} = M(1) \oplus M(\eta_k) \oplus \cdots \oplus M(\eta_k^{k-1})$$

where $M(\eta_k^j)$ denotes the complex matrix obtained by substituting $\eta_k^j = e^{2\pi i j/k}$ into M . Furthermore, any eigenvector of $h_*^{(k)}$ not lying in $S^{(k)}$ has eigenvalue which is a root of unity.

4 ENTROPY AND FIRST HOMOLOGY

The topological entropy, $h_{top}(f)$, is a well-known measure of the complexity of the dynamics of a self-map f of a compact metric space. See [AKM65], [DGS76] or [KH95] for more information. The next well-known lemma contains part of the main idea in Fried's paper [Fri86]: passing to a finite cover often allows one to detect more growth on homology.

Lemma 4.1 *If $f : Y \rightarrow Y$ is a continuous map of the compact manifold Y , then*

$$h_{top}(f) \geq \sup \left\{ \log \left(\text{sr} \left(\tilde{f}_* \right) \right) \right\}$$

where the supremum is over all lifts, $\tilde{f}$, of f to a finite cover $\tilde{Y}$ and $\tilde{f}_*$ is the action of $\tilde{f}$ on first homology of the cover $H_1(\tilde{Y}; \mathbb{R})$.

Proof: The lemma follows directly from two classic results. Manning proved in [Man74] that $h_{top}(f) \geq \log(\text{sr}(f_*))$ for any continuous $f : Y \rightarrow Y$ and Bowen proved in [Bow71] that entropy is preserved under finite to one factors, in particular, $h_{top}(f) = h_{top}(\tilde{f})$ for any lift $\tilde{f}$ of f to a finite cover. ■

For a braid β we define $h_{top}(\beta) = \inf \{ h_{top}(h) : h \in \beta \}$. Now Theorem 3.4 says that the maximal spectral radius of $B(e^{2\pi i j/k})$ for $0 \leq j < k$ gives the spectral radius on the first homology of the lift to the k -fold Burau cover $D^{(k)}$ of an $h \in \beta$. Thus by Lemma 4.1 we have what was referred to in the Introduction as the *Burau estimate*.

$$h_{top}(\beta) \geq \sup \{ \log(\text{sr}(B(e^{2\pi i j/k}))) : j/k \in \mathbb{Q} \}.$$

Since the entries in $B(t)$ are polynomials, $\text{sr}(B(t))$ is continuous in t . Thus we have the result of Fried [Fri86] and Kolev [Kol89]:

Lemma 4.2 *If $\beta \in B_n$ with Burau representation $B(t)$, then*

$$h_{top}(\beta) \geq \sup \{ \log(\text{sr}(B(\eta))) : \eta \in S^1 \}. \quad (4.1)$$

A fundamental result in Nielsen-Thurston theory says that $h_{top}(\beta) = h_{top}(\Phi)$ where Φ as given in Theorem 2.1 is the Thurston representative in the isotopy class β . A natural question is whether this value is detected by the action on homology in some finite cover, or in the current context, whether the estimate in (4.1) is ever sharp.

To investigate this question we must first understand a simpler question, namely, for a pseudo-Anosov map ϕ of a surface X , when is $h_{\text{top}}(\phi) = \text{sr}(\phi_*)$? In other words, when is Manning's estimate in [Man74] sharp for a pseudo-Anosov map? The answer turns out to depend exactly on the orientability of the ϕ -invariant foliations. It is well-known that the oriented, measured foliation $\mathcal{F}^u$ gives rise to a homology class $v \in H_1(X; \mathbb{R})$, for example, as an asymptotic direction as in Schwartzman ([Sch57]) or a geometric current as in Ruelle and Sullivan([RS75]). Briefly, v is the average direction in homology obtained from flowing along the one-dimensional leaves of $\mathcal{F}^u$. This average exists and is unique because the unstable foliation of a pseudo-Anosov map is uniquely ergodic ([FLP79]).

Now if the unstable foliation $\mathcal{F}^u$ gives rise to $v^u \in H_1(X; \mathbb{R})$, then since $\phi_*\mathcal{F}^u = \lambda\mathcal{F}^u$, on first homology we have $\phi_*v^u = \lambda v^u$. Thus v^u is an eigenvector of ϕ_* with eigenvalue λ , and so $\text{sr}(\phi_*) \geq \lambda$. On the other hand, since pseudo-Anosov maps have entropy equal to the logarithm of their expansion constant, Lemma 4.1 yields that $\lambda \geq \text{sr}(\phi_*)$. Thus when the invariant foliations are orientable we see that the spectral radius on first homology gives the entropy of a pseudo-Anosov map. The converse of this fact doesn't seem as well-known so we include a proof below for completeness.

Lemma 4.3 *Suppose $\phi : X \rightarrow X$ is a pseudo-Anosov homeomorphism of the orientable surface X having ℓ connected components (which, according to the definition of pseudo-Anosov map, must be permuted cyclically by ϕ), and let λ be the expansion constant of ϕ .*

- (a) *The pA map ϕ has orientable invariant foliations in some (hence each) connected component if and only if $\text{sr}(\phi_*) = \lambda$.*
- (b) *Suppose ϕ has oriented invariant foliations, and let $\epsilon = \pm 1$ according to whether ϕ preserves or reverses the orientation of the unstable foliation. Then each complex number of the form $\epsilon e^{2\pi i j / \ell} \lambda$ is a simple eigenvalue of ϕ_* , and every other eigenvalue μ satisfies $|\mu| < \lambda$.*

Proof: We first suppose that X is connected, i.e. that $\ell = 1$, and prove (b) and then (a). The proof when $\ell > 1$ is an easy modification.

When $\ell = 1$, (b) states that if the invariant foliations of ϕ are oriented, then $\epsilon\lambda$ is a simple eigenvalue of ϕ_* , and every other eigenvalue is smaller in modulus. Because X is orientable, orientability of the unstable foliation of ϕ is equivalent to orientability of the stable foliation of ϕ ([CLN85]), so we need only talk about orientability of the former.

In [Ryk99] Rykken shows that if the pseudo-Anosov map ϕ has an orientable unstable foliation, then except for zeros and roots of unity, the eigenvalues of its action on first homology are the same as those of its Markov transition matrix, $A_{\mathcal{P}}$, with respect to any Markov partition $\mathcal{P}$. Using this result (b) then follows from the Perron-Frobenius Theorem applied to $A_{\mathcal{P}}$ (see, for example, [Kit98]).

We noted above the theorem how one implication in (a) follows from treating the foliation as an asymptotic cycle. We prove the contrapositive of the converse and so assume that the unstable foliation $\mathcal{F}^u$ of ϕ is not orientable. Let $\vec{X} \xrightarrow{\vec{p}} X$ be the orientation double-cover of $\mathcal{F}^u$ constructed in Section 2.4; ϕ lifts to a pseudo-Anosov $\vec{\phi}$ of $\vec{X}$ with unstable foliation denoted $\vec{\mathcal{F}}^u$. The leaves of $\vec{\mathcal{F}}^u$ of necessity project under $\vec{p}$ to those of $\mathcal{F}^u$. Since $\vec{\phi}$ also

has expansion constant λ , by part (b) $\vec{\phi}_* : H_1(\vec{X}) \rightarrow H_1(\vec{X})$ has a simple eigenvalue equal to $\epsilon\lambda$ and all its other eigenvalues have smaller modulus.

If we let θ denote the generator of the deck group of $\vec{X}$, then θ will take leaves of $\vec{\mathcal{F}}^u$ to leaves reversing the orientation while preserving the transverse measure, and so $\theta_* v^u = -v^u$. Now since $\theta^2 = \text{id}$, $H_1(\vec{X}, \mathbb{R}) = V_+ \oplus V_-$ with $V_{\pm}$ the eigenspace of θ_* with eigenvalue ± 1 . Moreover, we now show that V_- is precisely the kernel of the map induced on homology by the covering projection $\vec{p}_* : H_1(\vec{X}, \mathbb{R}) \rightarrow H_1(X, \mathbb{R})$. Suppose τ is a 1-cycle in $\vec{X}$ avoiding the singular points of $\vec{\mathcal{F}}$ such that $\vec{p}_*[\tau] = 0 \in H_1(X)$. Then we may write $\vec{p}_*\tau = \partial\Delta$ where $\Delta = \sum_i a_i \Delta_i$ is some 2-chain. Lift each Δ_i to a 2-simplex Δ'_i in $\vec{X}$ and let $\vec{\Delta} = \sum_i a_i (\Delta'_i + \theta\Delta'_i)$. Clearly, $\partial\vec{\Delta} = \tau + \theta\tau$ proving that $[\tau] + [\theta\tau] = 0 \in H_1(\vec{X})$. In other words $[\tau] \in V_-$. So $\ker \vec{p}_* \subset V_-$. Conversely if $v \in V_-$, then $2\vec{p}_*(v) = \vec{p}_*(v) + \vec{p}_*(\theta(v)) = \vec{p}_*(v) - \vec{p}_*(v) = 0$, and hence $V_- = \ker \vec{p}_*$ as claimed.

Thus ϕ_* acting on $H_1(X, \mathbb{R})$ is conjugate to $\vec{\phi}_*$ acting on V_+ . Since $v^u \in V_-$ and the eigenvalue $\epsilon\lambda$ is geometrically simple, it follows that every eigenvalue of $\vec{\phi}_*$ having an eigenvector lying in V_+ has modulus strictly smaller than λ , so that $\text{sr}(\phi_*) < \lambda$ as required. ■

5 THE BURAU ESTIMATE FOR PSEUDO-ANOSOV BRAIDS

A braid $\beta \in B_n$ is said to be *pseudo-Anosov* if in the Thurston normal form $\Phi \in \beta$ of Theorem 2.1 the set A consists of just one annulus a which is the collar of ∂D , and the restriction of Φ to the complement of a is a pseudo-Anosov map ϕ . In this case we will call Φ a *collared pseudo-Anosov* map and consider its invariant foliations to be those of ϕ .

For simplicity of notation we let $r(\theta) = \text{sr}(B(e^{2\pi i\theta}))$ and just consider $\theta \in [0, 2\pi)$. If β is pseudo-Anosov with expansion constant $\lambda = e^{h_{\text{top}}(\Phi)}$, Lemma 4.1 shows that $r(\theta) \leq \lambda$ for all $\theta \in [0, 2\pi)$. In this section we investigate when the Burau estimate (4.1) is sharp for a pseudo-Anosov braid and the substitution of a root of unity, that is, when do we have $r(j/k) = \lambda$ for some $j/k \in \mathbb{Q}$. The first observation is that according to Theorem 3.4 when $r(j/k) = \lambda$ occurs, then the action of $h_*^{(k)}$ on the first homology of the k -fold Burau cover $H_1(D^{(k)})$ has some eigenvalue of modulus λ . By Lemma 4.3, this eigenvalue must be either λ or $-\lambda$, it must be simple, and all other eigenvalues of $h_*^{(k)}$ must be of smaller modulus. So, in particular, $r(\ell/k) < \lambda$ for all $\ell \neq j$.

The second observation is that the function $r(\theta)$ is periodic with period one and in addition, since the coefficients of the Laurent polynomial entries of $B(t)$ are real, $\text{sr}(B(\eta)) = \text{sr}(B(\bar{\eta}))$, where $\bar{\eta}$ denotes the complex conjugate of η . Thus r is an even, one-periodic function and so it is even about $1/2$.

Putting the two observations together we have that for a given k , $r(j/k) = \lambda$ for at most one j and since $r(1/2 + x) = r(1/2 - x)$, that can only happen if $j/k = 1/2$. Thus if k is even, the only possibility for the Burau estimate to be sharp is that $r((k/2)/k) = \lambda$ and so $r(j/k) < \lambda$ for other j . On the other hand, if k odd, $r(j/k) < \lambda$ for all j , so the Burau estimate is never sharp for substitutions with k odd.

In the next Proposition we connect these observations with the structure of the invariant foliations from Section 2.4 and Lemma 4.3.

Theorem 5.1 *Suppose $\beta \in B_n$ is a braid represented by the collared pseudo-Anosov map $\Phi : D \rightarrow D$ having expansion factor $\lambda > 1$. The following are equivalent:*

- (a) $\text{sr}(B(e^{2\pi ij/k})) = \lambda$ for some $k > 0$ and some $0 \leq j < k$;
- (b) $\text{sr}(B(-1)) = \lambda$ and -1 is the only root of unity for which this occurs.
- (c) The invariant foliations $\mathcal{F}^u$ and $\mathcal{F}^s$ have an odd-order singularity at each puncture of D , and all singularities of $\mathcal{F}^u$ and $\mathcal{F}^s$ in the interior of D have even order.
- (d) $D^{(2)}$ is the orientation double-cover of $\mathcal{F}^u$ and $\mathcal{F}^s$ (after removing the collaring).

Proof: The observations above the Theorem prove the equivalence of (a) and (b).

Let $\Phi^{(2)}$ be the preferred lift of Φ to $D^{(2)}$ and $\Phi_*^{(2)}$ its action on $H_1(D^{(2)})$. By Theorem 3.4, the eigenvalues of $\Phi_*^{(2)}$ are those of $B(1) \oplus B(-1)$, together with some roots of unity. Since $B(1)$ is a permutation, λ (or $-\lambda$) is an eigenvalue of $\Phi_*^{(2)}$ if and only if it is an eigenvalue of $B(-1)$. In addition, by Lemma 4.3, since $\Phi^{(2)}$ is collared pseudo-Anosov, one of λ or $-\lambda$ is an eigenvalue of $\Phi_*^{(2)}$ if and only if the invariant foliations of $\Phi^{(2)}$ are oriented. Thus (b) holds exactly when the lift of $\mathcal{F}^u$ to $D^{(2)}$ is oriented, and so in particular (d) implies (b).

As remarked above Lemma 2.2, if γ is a small loop surrounding a singularity P , then recalling that $\vec{\rho}$ denotes the epimorphism defining the orientation double cover, we have $\vec{\rho}([\gamma]) = 1$ iff P has odd order. On the other hand, the epimorphism $\tau_2 : H_1(D) \rightarrow \mathbb{Z}_2$ which yields the two-fold Burau cover $D^{(2)}$ is defined by $\tau_2([\gamma']) = 1$ for γ' a small loop around a puncture. This shows the equivalence of (c) and (d).

Now we will show (b) implies (c) by proving its contrapositive. If $\mathcal{F}^u$ has an interior singularity of odd order, by Lemma 2.2 the lifted foliations to $D^{(2)}$ are not oriented. Now note that by Euler-Poincaré-Hopf formula (2.1), $\mathcal{F}^u$ always has a one-pronged singularity at some puncture x_i of D . If $\mathcal{F}^u$ has an even order singularity at some other puncture x_j , we consider the homotopy class $\alpha = \gamma_j^{-1}\gamma_i$. We then have $\vec{\rho}(\alpha) = 0 + 1 = 1$ and $\tau_2(\alpha) = -1 + 1 = 0$. Thus there can be no homomorphism δ with $\delta \circ \rho_2 = \vec{\rho}$, and so again by Lemma 2.2, the lifted foliations to $D^{(2)}$ are not oriented. Since we have just seen that (b) holds exactly when the lift of $\mathcal{F}^u$ to $D^{(2)}$ is oriented, we have that (b) implies (c). ■

Remark 5.2 Lemma 5 and Theorem 7 in [SKL02] contain a portion of the above Theorem in slightly different language, namely, the implications (c) $\implies$ (d) and (d) $\implies$ (b).

6 THE BURAU ESTIMATE FOR REDUCIBLE BRAIDS

6.1 PSEUDO-ANOSOV COMPONENTS AND BURAU ORIENTABILITY

The Thurston normal form $\Phi = \Phi_1 \cup \dots \cup \Phi_m$ of a general braid $\beta \in B_n$ can be quite complicated. Using the notation of Theorem 2.1, there may be several reducing annuli forming the set A , each component $\Phi_i = \Phi|_{X_i}$ of Φ may be either periodic or pseudo-Anosov, and X_i itself may have one or more connected components. In this section we will consider this more complicated decomposition and obtain general results on the sharpness of the entropy bound provided by the Burau representation.

The simplest case is when each Φ_i is periodic, and so $h_{top}(\Phi) = 0$. Since all the eigenvalues of Φ_* are roots of unity, we have $\text{sr}(B(1)) = 1$, giving a sharp bound on the entropy. Thus from now on we suppose that some component of Φ is pseudo-Anosov, and we let $\lambda = e^{h_{top}(\Phi)}$. The main theorem of this section gives necessary and sufficient conditions for a root of unity ω to make the Burau estimate (4.1) sharp, i.e. to satisfy $\text{sr}(B(\omega)) = \lambda$.

To state these conditions, we first define for each component Φ_i a positive integer a_i expressing the manner in which the surface X_i on which Φ_i is supported surrounds the punctures of D . Begin by choosing a component Φ_i of Φ . The supporting surface X_i of Φ_i may have several connected components; by definition, these are permuted cyclically by Φ_i . Let X_{i0} be one of these components. Thus X_{i0} is a finitely punctured subdisk of D from which a finite collection of open punctured subdisks have been deleted (see Figure 2(a)). Let $x_1, \dots, x_{r'}$ denote the punctures in X_{i0} , where $0 \leq r' \leq n$, and let $O_{r'+1}, \dots, O_r$ be the deleted subdisks. For $r' < j \leq r$ we write m_j for the number of punctures of D which lie in O_j ; and for $1 \leq j \leq r'$ we define $m_j = 1$. Finally, we define a_i by

$$a_i = \gcd(m_1, m_2, \dots, m_r). \quad (6.1)$$

Because Φ permutes the connected components of X_i cyclically, and sends punctures to punctures, this definition is independent of the choice of X_{i0} .

The next lemma gives another interpretation of the number a_i which follows from the definition of the morphism $\tau : H_1(D) \rightarrow \mathbb{Z}$ associated to the covering space $D^{(\infty)}$.

Lemma 6.1 *Let ℓ_i be the number of connected components of the supporting surface X_i of Φ_i , and let X_{i0} be the chosen connected component of X_i as above. Write $X_i^{(\infty)} = (p^{(\infty)})^{-1}(X_i)$ and $X_{i0}^{(\infty)} = (p^{(\infty)})^{-1}(X_{i0})$. The number a_i just defined is equal to the number of connected components of $X_{i0}^{(\infty)}$. Hence the number of connected components of $X_i^{(\infty)}$ is $\ell_i a_i$.*

Proof: According to Thurston's Theorem 2.1, the waist curve of any reducing annulus $a \in A$ is neither null-homotopic nor homotopic to a puncture. It follows that the inclusion $X_i \subset D$ induces an injection $H_1(X_i) \rightarrow H_1(D)$, and so we can consider $H_1(X_i)$ as a subgroup of $H_1(D)$. Then $X_i^{(\infty)} \rightarrow X_i$ is (isomorphic to) the covering space defined by the homomorphism $\tau|_{H_1(X_i)}$. By definition τ sends a small clockwise loop around the puncture x_j (respectively, the hole O_j) of X_{i0} to $m_j \in \mathbb{Z}$, and therefore a_i is the generator of $\text{im } \tau|_{H_1(X_{i0})}$. This proves the first statement of the lemma. Since h permutes the connected components of X_i cyclically and since $\tau\Phi_* = \tau$, we have that $X_i^{(\infty)}$ has the same number of connected components above each component of X_i , proving the second statement. ■

Let us say that a pseudo-Anosov component Φ_i of Φ is *Burau orientable* provided the lifts of its invariant foliations to some $D^{(k)}$ are orientable. The main theorem of this section is:

Theorem 6.2 *Let $\Phi = \Phi_1 \cup \dots \cup \Phi_m$ be the Thurston normal form of the braid β , and for each i such that Φ_i is pseudo-Anosov, let a_i be as in (6.1).*

1. *The pseudo-Anosov component Φ_i is Burau orientable if and only if (with notation as in the definition of a_i) the invariant foliations of Φ_i have a singularity of odd order at*




Figure 2: (a) A typical supporting surface X_i . Here $r' = 2$, $r = 3$, $m_3 = 2$ and $a_i = 1$. (b) the braid β_3 and its Thurston decomposition.

each puncture x_j of X_{i0} , and a singularity of odd (even) order on the boundary of each deleted disk O_j such that m_j/a_i is odd (even).

2. Let I be the set of $1 \leq i \leq m$ such that Φ_i is pseudo-Anosov, Burau orientable, and satisfies $h_{top}(\Phi_i) = h_{top}(\Phi)$. Then the set of roots of unity ω for which $\text{sr}(B(\omega)) = \lambda$, is equal to the union over $i \in I$ of the set of all a_i -th roots of -1 .

In particular, if I is empty, then $\text{sr}(B(\omega)) < \lambda$ for every root of unity ω .

Before embarking on the proof of Theorem 6.2, we will illustrate the theorem with some examples.

6.2 EXAMPLES

It is convenient to use the following notation. Let $n > 0$. If i, n_1, n_2 are positive integers with $i + n_1 + n_2 - 1 \leq n$, we let $\sigma_{i,n_1,n_2} \in B_n$ denote the braid which moves the group of n_1 consecutive strings starting at string i over the group of n_2 consecutive strings starting at string $i + n_1$:

$$\sigma_{i,n_1,n_2} = (\sigma_{i+n_1-1} \cdots \sigma_{i+n_1+n_2-2})(\sigma_{i+n_1-2} \cdots \sigma_{i+n_1+n_2-3}) \cdots (\sigma_i \cdots \sigma_{i+n_2-1}).$$

In particular $\sigma_{i,1,1} = \sigma_i$ for all $1 \leq i \leq n - 1$.

Example 6.3 Let $n' \geq 1$, and define a braid $\beta_{n'}$ on $3n'$ strings by setting

$$\beta_{n'} = \sigma_{1,n',n'} \sigma_{n'+1,n',n'}^{-1}.$$

Figure 2(b) shows $\beta_{n'}$ when $n' = 3$. The Thurston normal form Φ of $\beta_{n'}$ reduces along 4 annuli $A = A_0 \cup A_1 \cup A_2 \cup A_3$: there is one such annulus collaring the boundary of D , and one surrounding each of the three groups of n' punctures. The component of Φ in the outer connected component of $D \setminus A$ is pseudo-Anosov. Call this component Φ_1 and its supporting surface X_1 . The other component of Φ is periodic, and it cyclically permutes




Figure 3: The Burau estimate (a) for β_8 and (b) for β_5 . The horizontal lines represent the growth rate $\lambda \sim 2.618$.

the three inner connected components of $D \setminus A$. The Euler-Poincaré-Hopf formula (2.1) shows that the invariant foliations of Φ_1 have four 1-pronged singularities, one at each of the four boundary components of X_1 , and no other singularities. According to (6.1) we have $a_1 = m_1 = m_2 = m_3 = n'$, and hence Φ_1 is Burau orientable by the first part of Theorem 6.2.

The growth rate of $\beta_{n'}$ is $\lambda = e^{h_{top}(\Phi_1)} \sim 2.618$. Figure 3 shows the graph of the map sending $\theta \in [0, 1]$ to the spectral radius of the substituted Burau matrix $B(e^{2\pi i \theta})$ of $\beta_{n'}$, in the two cases $n' = 8 = 2^3$ and $n' = 5$.

Here is one part of the justification of Theorem 6.2 for this example. According to Lemma 6.6 below, if k is a multiple of $2n'$, the invariant foliations of Φ_1 lift to orientable foliations in the k -fold covering space $X_1^{(k)}$ of X_1 . Moreover this covering space has exactly n' connected components, each fixed by the lift of Φ . Therefore by Theorem 3.4 and Lemma 4.3 we expect the matrices $B(\omega)$, for ω a k -th root of unity, to contribute exactly n' eigenvalues equal to λ or $-\lambda$ (counted with multiplicity). Theorem 6.2 states, in addition, that it is precisely the n' -th roots of -1 which contribute these eigenvalues, a fact clearly reflected in Figure 3.

Example 6.4 Let $\beta'_1 \in B_9$ and $\beta'_2 \in B_8$ be the braids

$$\beta'_1 = \sigma_{1,3,3} \cdot \sigma_{4,3,3}^2 \cdot \sigma_{1,3,3}^3$$

and

$$\beta'_2 = \sigma_{1,3,3} \cdot \sigma_{4,3,2} \cdot \sigma_{4,2,3} \cdot \sigma_{1,3,3}^3.$$

Note that β'_1 respects the grouping of the punctures into consecutive groups of three, while β'_2 respects the grouping of the punctures into two consecutive groups of three and one of two.




Figure 4: The Burau estimate (a) for β'_1 and (b) for β'_2 . The horizontal lines represent the growth rate $\lambda \sim 5.828$.

As for the previous example, the Thurston normal form Φ of β'_i has one reducing annulus around each of the three groups of punctures. Each such annulus encloses a periodic (in fact fixed) component of Φ . If X_1 denotes the complement of the reducing annuli, then $\Phi_1 = \Phi|_{X_1}$ is pseudo-Anosov with growth rate $\lambda \sim 5.828$. Again the invariant foliations of Φ_1 have a one-pronged singularity on each boundary component of X_1 .

Now for β'_1 we have $a_1 = m_1 = m_2 = m_3 = 3$, and so Theorem 6.2 shows that Φ_1 is Burau orientable. Therefore the Burau estimate is sharp at all of the cubic roots of -1 (see Figure 4(a)). For β'_2 , however, we have $m_1 = m_2 = 3$ and $m_3 = 2$, and so $a_1 = \gcd(2, 3) = 1$. Thus the conditions of the first part of Theorem 6.2 fail to hold and Φ_1 is not Burau orientable, so the Burau estimate is never sharp (see Figure 4(b)).

Example 6.5 Let $\beta'' \in B_{10}$ be the braid

$$\beta'' = \beta_3 \cdot \sigma_{1,9,1} \cdot \sigma_{1,1,9}$$

where $\beta_3 \in B_9 \subset B_{10}$ is as in Example 6.3 above². See Figure 5(a). The Thurston normal form Φ of β'' has one pseudo-Anosov component $\Phi_1 = \Phi|_{X_1}$, topologically conjugate to the pseudo-Anosov component of the Thurston normal form of B_3 . By Theorem 6.2 this component is Burau orientable and the Burau estimate is sharp at each of the cubic roots of -1 . However, β'' is distinguished from β_3 by the fact that the eigenvalue of $B(-e^{2\pi ij/3})$ of modulus λ is not always real (see Figure 5(b)). This can be seen as follows. Lemma 6.6 below shows that for k a multiple of 6 the lift of Φ_1 to the k -fold cover $X_1^{(k)}$ has orientable foliations. Because Φ acts as a full twist in the outer component of D , the three connected components of $X_1^{(k)}$ are permuted cyclically by $\Phi^{(k)}$. So by Theorem 3.4 and Lemma 4.3, the matrices $B(\omega)$, for ω a k -th root of unity, contribute to $\Phi_*^{(k)}$ exactly three eigenvalues of

²As is usual we regard B_9 as the subgroup of B_{10} generated by the first eight generators $\sigma_1, \dots, \sigma_8$.




Figure 5: (a) The braid β'' ; (b) the eigenvalues of $B(e^{2\pi i\theta})$ for $\theta \in [0, 1]$. Here the rightmost axis represents the unit interval $[0, 1]$ and the other two axes represent the complex plane.

modulus λ , differing from each other by the cubic roots of unity. These eigenvalues are the extremal points of the curve in Figure 5(b).

6.3 PROOF OF THEOREM 6.2

We will prove Theorem 6.2 by a series of lemmas. The main idea of the proof, already suggested by the above examples, is the following. As we lift a pseudo-Anosov component Φ_i of the Thurston normal form Φ of the chosen braid β into successive covering spaces $D^{(k)}$, the supporting surface of the lift of Φ_i may become disconnected. If the lifted invariant foliations also become orientable, then Lemma 4.3 implies the existence of several eigenvalues of modulus λ (counted with geometric multiplicity). By Theorem 3.4 these eigenvalues are distributed among those of the matrices $B(\omega)$, where ω is an k -th root of unity. An analysis of the action of $\Phi^{(\infty)}$ on the first homology of $X_i^{(\infty)}$ then shows that it is precisely the a_i -th roots of -1 which contribute such an eigenvalue.

We begin with a lemma which provides a stronger version of the first part of Theorem 6.2. Let $\Phi = \Phi_1 \cup \dots \cup \Phi_m$ be the Thurston normal form of the braid $\beta \in B_n$, let Φ_i be a pseudo-Anosov component of Φ , and let $\mathcal{F}^u$ and $\mathcal{F}^s$ denote the invariant foliations of Φ_i . For $k \geq 1$ we consider the k -fold Burau covering space $p^{(k)} : D^{(k)} \rightarrow D$ and write $X_i^{(k)} = (p^{(k)})^{-1}(X_i)$ and $\Phi_i^{(k)} = \Phi^{(k)}|_{X_i^{(k)}}$. As in the definition of a_i in (6.1), let X_{i0} be a connected component of X_i , and let $x_1, \dots, x_{r'}$ be the punctures in X_{i0} and $O_{r'+1}, \dots, O_r$ the deleted disks. Further, for each $1 \leq j \leq r$ let κ_j be the order of the singularity which $\mathcal{F}^u$ and $\mathcal{F}^s$ exhibit at x_j (if $j \leq r'$) or on ∂O_j (if $r' < j \leq r$). We have:

Lemma 6.6 *Let $\Phi_i = \Phi|_{X_i}$ be a pseudo-Anosov component of the Thurston normal form Φ of β , and let a_i be defined as in (6.1). Write $a = a_i$ and suppose that $a = 2^u a'$ where a' is*

odd. Then the following are equivalent:

- (1) Φ_i is Burau orientable, i.e. there exists $k \geq 1$ such that the lifts of $\mathcal{F}^u$ and $\mathcal{F}^s$ to $X_i^{(k)}$ are orientable;
- (2) The set of k for which the lifts of $\mathcal{F}^u$ and $\mathcal{F}^s$ to $X_i^{(k)}$ are orientable is the set of multiples of 2^{u+1} .
- (3) All singularities of $\mathcal{F}^u$ and $\mathcal{F}^s$ in the interior of X_{i0} have even order, and for all $1 \leq j \leq r$, $m_j/a \equiv \kappa_j \pmod{2}$. In other words m_j is an odd (even) multiple of a whenever κ_j is odd (even).

Proof: The implication (2) $\implies$ (1) is trivial; we will prove (1) $\implies$ (3) and (3) $\implies$ (2). Except for an application of Lemma 2.2, the proof is essentially algebraic – it involves only the relevant morphisms of $H_1(X_i)$ into $\mathbb{Z}$ and $\mathbb{Z}_k$. For $1 \leq j \leq r$ let α_j be the homology class of a small clockwise loop around the puncture x_j (if $j \leq r'$) or around the disk O_j (if $j > r'$). Then $H_1(X_{i0})$ can be identified with the subgroup of $H_1(D)$ generated by the α_j 's. By definition $\tau(\alpha_j) = m_j$ for each j , and $\tau_k(\alpha_j) = \xi_k \tau(\alpha_j) = m_j \pmod{k}$, where $\xi_k : \mathbb{Z} \rightarrow \mathbb{Z}_k$ is the quotient homomorphism. When all singularities of $\mathcal{F}^u$ and $\mathcal{F}^s$ in the interior of X_i have even order, we let $\vec{\rho} : H_1(X_i) \rightarrow \mathbb{Z}_2$ be the morphism associated to the orientation cover of $\mathcal{F}^u$ and $\mathcal{F}^s$. Because Φ_i preserves the foliations and permutes the components cyclically, $\vec{\rho}$ is determined by its values on $H_1(X_{i0})$: namely $\vec{\rho}(\alpha_j) = \kappa_j \pmod{2}$ for all j .

According to Lemma 2.2 we know that for $k \geq 1$ the following two statements are equivalent:

- (i) The lifts of $\mathcal{F}^u$ and $\mathcal{F}^s$ to $X_i^{(k)}$ are orientable;
- (ii) All singularities in the interior of X_i have even order, and there exists a homomorphism $\delta_k : \text{im } \tau_k|_{H_1(X_i)} \rightarrow \mathbb{Z}_2$ such that $\delta_k \circ \tau_k = \vec{\rho}$ on $H_1(X_i)$.

In addition, since $\tau_k \Phi_* = \tau_k$, the formula $\delta_k \circ \tau_k = \vec{\rho}$ holds on $H_1(X_i)$ whenever it holds on $H_1(X_{i0})$.

We note some elementary facts about cyclic groups. For $q \geq 1$ odd, there is no nontrivial homomorphism $\mathbb{Z}_q \mapsto \mathbb{Z}_2$, while for q even there is only one such homomorphism: namely, the homomorphism which sends odd multiples of the generator to 1 and even multiples to 0 (a property which is independent of the choice of generator). And if q is odd then every subgroup of $\mathbb{Z}_q$ has odd order.

Suppose that (1) holds, that is, that Φ_i is pseudo-Anosov and for some $k > 0$ the lifts of $\mathcal{F}^u$ and $\mathcal{F}^s$ to $X_i^{(k)}$ are orientable. Since $p^{(k)} : X_i^{(k)} \rightarrow X_i$ is ramified only around the punctures, this implies that all singularities in the interior of X_i have even order. Since $\text{im } \tau|_{H_1(X_i)} = a\mathbb{Z}$, we have $\text{im } \tau_k|_{H_1(X_i)} = \xi_k(a\mathbb{Z}) = \xi_k(a)\mathbb{Z}_k$. The Euler-Poincare-Hopf formula (2.1) shows that $\mathcal{F}^u$ and $\mathcal{F}^s$ have at least one 1-pronged singularity in X_{i0} , which must lie at a puncture or on the boundary of a deleted disk. Thus $\vec{\rho}$ is onto, so from the previous paragraph we see that (ii) holds precisely when k is even and for all $1 \leq j \leq r$ we have: $\tau_k(\alpha_j)$ is an odd (even) multiple of $\xi_k(a)$ iff κ_j is odd (even). If k is even then ξ_k sends odd (even) multiples of a to odd (even) multiples of $\xi_k(a)$ and so this implies the same statement in $\mathbb{Z}$: that is, that (3) holds. Thus (1) $\implies$ (3).

On the other hand suppose (3) holds and set $k = 2^{u+1}$. Since $a = 2^u a'$ where a' is odd, we see that $\text{im } \tau_k|_{H_1(X_i)}$ is the subgroup $\{0, \xi_k(a)\}$ of $\mathbb{Z}_k$ having order two. Let δ_k be the isomorphism of this subgroup onto $\mathbb{Z}_2$. Again, since k is even, to say that $\xi_k(m_j) = \xi_k(a)$ is to say that m_j is an odd multiple of a . By (3) this happens precisely when κ_j is odd; hence $\delta_k \tau_k = \vec{\rho}$ on $H_1(X_i)$. Therefore since (i) $\Leftrightarrow$ (ii) the lifts of $\mathcal{F}^u$ and $\mathcal{F}^s$ to $X_i^{(k)}$ are orientable. Therefore the set of k for which the lifted foliations in $X^{(k)}$ are orientable contains 2^{u+1} , and, it follows, also all multiples of 2^{u+1} .

To complete the proof that (3) $\implies$ (2), suppose for a contradiction that (3) holds and that for some k which is not a multiple of 2^{u+1} , the lifts of $\mathcal{F}^u$ and $\mathcal{F}^s$ to $X_i^{(k)}$ are orientable. The last condition implies that k is even, so we must have $u > 0$. Let δ_k be the homomorphism supplied by (ii). Since $\vec{\rho}$ is onto, δ_k must send the generator $\xi_k(a)$ of $\text{im } \tau_k|_{H_1(X_i)}$ to $1 \in \mathbb{Z}_2$. Now write $k = 2^{u'} k'$ where k' is odd and $u' \leq u$. The order of $\xi_k(a)$ in $\mathbb{Z}_k$ is given as $\zeta = \text{lcm}(a, k)/a = \text{lcm}(a', k')/a'$, which is odd. But then $0 = \delta_k(0) = \delta_k(\zeta \xi_k(a)) = \zeta \delta_k \xi_k(a) = \delta_k \xi_k(a) = 1$, a contradiction. Therefore no such k can exist. This completes the proof. ■

To prove the second part of Theorem 6.2, we will require two more lemmas. The first of these expresses $H_1(D^{(\infty)})$ in terms of the first homology groups of the $X_i^{(\infty)}$'s. Since each $X_i^{(\infty)}$ is $\Phi^{(\infty)}$ -invariant, we will be able to use this lemma to factorize the characteristic polynomial of the reduced Burau matrix $B(\beta)$.

Lemma 6.7 *Let $\Phi = \Phi_1 \cup \dots \cup \Phi_m \in \beta$ be as above and let $\Phi^{(\infty)}$ be the lift of Φ to $D^{(\infty)}$. Let $X_i^{(\infty)} = (p^{(\infty)})^{-1}(X_i)$ be the pull-back to $D^{(\infty)}$ of the supporting surface X_i of Φ_i . Then, as an Abelian group, $H_1(D^{(\infty)})$ decomposes as a direct sum of subgroups*

$$H_1(D^{(\infty)}) = \left(\bigoplus_{i=1}^m H_1(X_i^{(\infty)}) \right) \oplus V \quad (6.2)$$

in which each $H_1(X_i^{(\infty)})$ is $\Phi_*^{(\infty)}$ -invariant, and V is a free abelian group of finite rank.

Proof: We use the Mayer-Vietoris sequence inductively. Let $A^{(\infty)} = (p^{(\infty)})^{-1}(A)$ be the set of points of $D^{(\infty)}$ covering the reducing annuli in D . For $1 \leq i \leq m$ define A_i to be the set of those connected components of $A^{(\infty)}$ whose boundary intersects $X_i^{(\infty)}$, but does not intersect $X_j^{(\infty)}$ for any $1 \leq j < i$. Then for each i we let $Z_i = X_i^{(\infty)} \cup A_i$. Also, for $0 \leq i \leq m$ let $Y'_i = \bigcup_{j>i} Z_j$ and when $i > 0$ let $Y_i = Y'_i \cup A_i$. These spaces satisfy

$$Z_i \cap Y_i = A_i, \quad Z_i \cup Y_i = Y'_{i-1}$$

for each $1 \leq i \leq m$, and

$$Y'_0 = D^{(\infty)}, \quad A_m = Y'_m = Y_m = \emptyset.$$

Since $X_i^{(\infty)}$ is $\Phi^{(\infty)}$ -invariant for each i , so too are A_i, Z_i, Y_i and Y'_i .

For any reducing annulus $a \in A$, each boundary component of a is also the boundary of some (possibly punctured) subdisk D' of D . In fact, since no component of ∂a is allowed to

be null-homotopic, D' must have at least one puncture and so we will have $\tau([\partial D']) \neq 0$. It follows that any connected component of $A^{(\infty)}$ covering a is an infinite strip, isomorphic as a covering space to the universal cover of a . In particular $H_1(A_i) = 0$ for each i .

Therefore, regarding Z_i and Y_i as subspaces of their union Y'_{i-1} , the Mayer-Vietoris sequence for the pair (Z_i, Y_i) contains the segment

$$0 \longrightarrow H_1(Z_i) \oplus H_1(Y_i) \longrightarrow H_1(Y'_{i-1}) \xrightarrow{\partial} H_0(A_i) \longrightarrow \cdots \quad (6.3)$$

All groups in the above sequence are free abelian. Noting that $X_i^{(\infty)}$ is a deformation retract of Z_i and that Y'_{i-1} is a deformation retract of Y_{i-1} , we have

$$H_1(Y_{i-1}) \cong H_1(X_i^{(\infty)}) \oplus H_1(Y_i) \oplus V_i$$

where V_i is canonically isomorphic to $\text{im } \partial \subset H_0(A_i)$. Since A_i has finitely many connected components, $H_0(A_i)$ and V_i each have finite rank. The lemma is proved by applying this formula inductively, starting with $H_1(Y_{m-1}) = H_1(X_m^{(\infty)})$, and setting $V = \bigoplus_{i=1}^m V_i$. ■

Corollary 6.8 *For each $1 \leq i \leq m$ let $\Phi_i^{(\infty)}$ be the restriction of $\Phi^{(\infty)}$ to $X_i^{(\infty)}$, and denote by g_i the action of $\Phi_i^{(\infty)}$ on $H_1(X_i^{(\infty)})$. As in Section 3, regard $H_1(D^{(\infty)})$ and $H_1(X_i^{(\infty)})$ as modules over the ring $R = \mathbb{Z}[t^{\pm 1}]$. Then $\bigoplus_{i=1}^m H_1(X_i^{(\infty)})$ is a submodule of $H_1(D^{(\infty)})$ of full rank. Moreover, the characteristic polynomial of the module isomorphism $h_*^{(\infty)} : H_1(D^{(\infty)}) \rightarrow H_1(D^{(\infty)})$ is given by*

$$\chi(h_*^{(\infty)}) = \prod_{i=1}^m \chi(g_i).$$

Proof: Let $G = \bigoplus_{i=1}^m H_1(X_i^{(\infty)})$. Because the R -module structure on $H_1(D^{(\infty)})$ and on $H_1(X_i^{(\infty)})$ are each defined using the same group of deck transformations acting respectively on $D^{(\infty)}$ and on $X_i^{(\infty)}$, they coincide, making G a submodule of $H_1(D^{(\infty)})$.

As an R -module, $H_1(D^{(\infty)})$ is free and of rank $n - 1$. We claim that $G \subseteq H_1(D^{(\infty)})$ also has rank $n - 1$. For suppose $\text{rank}(G) < n - 1$. We may choose $w \in H_1(D^{(\infty)})$ so that $G \cup \{w\}$ spans a submodule of $H_1(D^{(\infty)})$ of rank strictly larger than $\text{rank}(G)$. Since $H_1(D^{(\infty)})$ is free, it follows that $w + G$ generates a free R -submodule of $H_1(D^{(\infty)})/G$. But in the notation of Lemma 6.7, we have $H_1(D^{(\infty)})/G \cong V$ as Abelian groups. Therefore $H_1(D^{(\infty)})/G$ has finite rank as an Abelian group, a contradiction.

By definition $h_*^{(\infty)}|_G = \bigoplus_{i=1}^m g_i$. Because G is a submodule of $H_1(D^{(\infty)})$, the characteristic polynomial of $h_*^{(\infty)}|_G$ divides the characteristic polynomial of $h_*^{(\infty)}$ on $H_1(D^{(\infty)})$. Since G has full rank both polynomials have the same leading coefficient x^{n-1} and so they are equal. This completes the proof. ■

The next lemma shows how the characteristic polynomial $\chi(g_i)$ of g_i reflects the permutation induced by $\Phi_i^{(\infty)}$ on the set of connected components of $X_i^{(\infty)}$.

Lemma 6.9 *Let $\Phi_i = \Phi|_{X_i}$ be a component of the Thurston normal form h of β , and let $\Phi_i^{(\infty)}$ be the lift of Φ_i to $X_i^{(\infty)}$. Let g_i denote the action of $\Phi_i^{(\infty)}$ on $H_1(X_i^{(\infty)})$. Suppose X_i*

has ℓ_i connected components, and let $d \in \mathbb{Z}$ be chosen so that $(\Phi^{(\infty)})^{\ell_i}(Y) = T^d(Y)$ for any connected component Y of $X_i^{(\infty)}$. Then there exists an integer $e > 0$ such that the characteristic polynomial of g_i is of the form

$$\chi(g_i) = x^{\ell_i e} + t^d P_1(t) x^{\ell_i(e-1)} + t^{2d} P_2(t) x^{\ell_i(e-2)} + \dots + t^{ed} P_e(t) \quad (6.4)$$

where each $P_j \in \mathbb{Z}[t^{\pm a_i}]$.

Proof: As before, we let X_{i0} be a connected component of X_i and $X_{i0}^{(\infty)}$ its pull-back to $D^{(\infty)}$. Let e be the dimension of $H_1(X_{i0}^{(\infty)})$ as a module over $\mathbb{Z}[t^{\pm 1}]$. Let $\{v_j : 1 \leq j \leq e\}$ be a basis for $H_1(X_{i0}^{(\infty)})$. We choose this basis so that all of the v_j 's are represented by loops in one and the same connected component, Y say, of $X_{i0}^{(\infty)}$. Pushing forward this basis under each of the first ℓ_i iterates of g_i , we obtain a basis for $H_1(X_i^{(\infty)})$. With respect to this basis, the matrix of g_i has block form

$$g_i = \begin{pmatrix} 0 & & \dots & \Omega \\ \text{id} & 0 & & \\ & \text{id} & 0 & \vdots \\ & & \ddots & \ddots \\ & & & \text{id} & 0 \end{pmatrix},$$

having $\ell_i \times \ell_i$ blocks each of dimension $e \times e$, where Ω is some matrix in $\text{GL}(e, \mathbb{Z}[t^{\pm 1}])$.

The connected components of $X_{i0}^{(\infty)}$ are in one to one correspondence with the cosets of $a_i \mathbb{Z}$ in $\mathbb{Z}$. Now $(\Phi^{(\infty)})^{\ell_i}$ sends the chosen connected component $Y \subset X_{i0}^{(\infty)}$ to the possibly different connected component $T^d Y$ of $X_{i0}^{(\infty)}$, where $d \in \mathbb{Z}$ is well-defined up to adding a multiple of a_i . It follows that the entries of Ω lie in the coset $t^d \mathbb{Z}[t^{\pm a_i}]$ of $\mathbb{Z}[t^{\pm a_i}]$ in $\mathbb{Z}[t^{\pm 1}]$.

Any product of j elements of Ω therefore lies in the coset $t^{jd} \mathbb{Z}[t^{\pm a_i}]$ of $\mathbb{Z}[t^{\pm a_i}]$. The lemma is proved when we note that the characteristic polynomial of the matrix of g_i is of the form

$$x^{\ell_i e} + \omega_1 x^{\ell_i(e-1)} + \dots + \omega_e$$

where for each $1 \leq j \leq e$, ω_j is a sum of products of j elements of Ω , so that $\omega_j \in t^{jd} \mathbb{Z}[t^{\pm a_i}]$. ■

Corollary 6.10 *The roots of $\chi(g_i)$ are naturally divided into sets each of $\ell_i a_i$ roots. If (x, t) and (x', t') are two roots of $\chi(g_i)$ in the same set, then $(x', t') = (\mu x, \nu t)$, where ν is some a_i -th root of unity and μ is some ℓ_i -th root of ν^d .*

Proof: Let $(x, t) \in \mathbb{C} \times \mathbb{C}$ and let μ and ν be as in the statement. A quick calculation using (6.4) shows that the value of $\chi(g_i)$ at $(\mu x, \nu t)$ is exactly ν^{ed} times its value at (x, t) . In particular if (x, t) is a root so too is $(\mu x, \nu t)$. ■

Proof of Theorem 6.2: Lemma 6.6 establishes the first part of Theorem 6.2. To prove the second part, recall that I is the set of $1 \leq i \leq m$ such that Φ_i is pseudo-Anosov, Burau

orientable, and satisfies $h_{top}(\Phi_i) = h_{top}(\Phi)$. For $k > 0$ we write $\eta_k = e^{2\pi i/k}$. Also, for each i we write $\Phi_i^{(k)} = \Phi^{(k)}|_{X_i^{(k)}}$ and $g_i^{(k)} = \Phi_*^{(k)}|_{H_1(X_i^{(k)})}$ as above.

Let $i \in I$. With notation as above, the number of connected components of $X_i^{(\infty)}$ is $\ell_i a_i$. Since $\tau_k = \xi_k \tau$, the same statement is true if we replace $X_i^{(\infty)}$ by $X_i^{(k)}$, for any k which is a multiple of a_i .

Suppose in fact that k is a multiple of $2a_i$. By Lemma 6.6, the invariant foliations of $\Phi_i^{(k)}$ in $X_i^{(k)}$ are orientable. The $\ell_i a_i$ connected components of $X_i^{(k)}$ are permuted by $\Phi_i^{(k)}$ in cycles all having the same number of components. Suppose the number of such cycles is L , so that each cycle contains $\ell_i a_i / L$ components. Lemma 4.3 applies to each such cycle, and we conclude that $g_i^{(k)}$ has exactly $\ell_i a_i / L$ eigenvalues of modulus λ , each having geometric multiplicity L . Indeed $\epsilon \lambda$ is one such eigenvalue, where $\epsilon \in \{1, -1\}$ is chosen according to whether $\Phi_i^{(k)}$ preserves or reverses the orientation of its unstable foliation. Counted with geometric multiplicity, there are exactly $\ell_i a_i$ eigenvalues of modulus λ .

Theorem 3.4 now implies that these $\ell_i a_i$ eigenvalues are distributed among the eigenvalues of the matrices $M(\eta_k^j)$, where M denotes the matrix of $g_i : H_1(X_i^{(\infty)}) \rightarrow H_1(X_i^{(\infty)})$ as in Section 3. Because $M(\nu_k^j)$ is obtained by substituting ν_k^j into M , an eigenvalue x of $M(\eta_k^j)$ corresponds to a root of $\chi(g_i)$ of the form (x, ν_k^j) . In particular, setting $k = 2a_i$, we conclude that there is some $0 \leq j_0 < 2a_i$ such that $(\epsilon \lambda, \eta_{2a_i}^{j_0})$ is a root of $\chi(g_i)$.

Meanwhile, since by Lemma 6.6 the lifts of $\mathcal{F}^u$ and $\mathcal{F}^s$ to $X_i^{(a_i)}$ are not orientable, a similar argument shows that no a_i -th root of unity can occur in such a root of $\chi(g_i)$. Therefore j_0 must be odd. By Corollary 6.10 we now see that every element of the set

$$J_i = \{(\mu \epsilon \lambda, \eta_{2a_i}^{j_0+j}) | j \text{ is even and } \mu \text{ is an } \ell_i\text{-th root of } \eta_{2a_i}^{jd}\}$$

is a root of $\chi(g_i)$, where d is as in Lemma 6.9. Since j_0 is odd, the roots of unity occurring in elements of J_i are precisely the a_i -th roots of -1 .

For k a multiple of $2a_i$, again by Theorem 3.4 we see that J_i accounts for all of the $\ell_i a_i$ eigenvalues of $g_i^{(k)}$ of modulus λ . Furthermore, every root of unity can be written as a k -th root of unity for some k which is a multiple of $2a_i$. It follows that J_i is precisely the set of roots (x, ν) of $\chi(g_i)$ such that ν is a root of unity and $|x| = \lambda$.

By Corollary 6.8, the characteristic polynomial of $B(\beta)$ is the product of those of the g_i 's. Therefore the set of roots of $\chi(B(\beta))$ is the union of those of the g_i 's. For $i \in I$, we have just accounted for all of the roots (x, ν) of $\chi(g_i)$ with $|x| = \lambda$ and ν a root of unity. For $i \notin I$ a similar argument shows that $\chi(g_i)$ can have no such roots. This completes the proof. ■

Proof of Theorem 1.2: Fix $\beta \in B_n$ and let $\Phi = \Phi_1 \cup \dots \cup \Phi_m$ be its Thurston normal form. If the Burau estimate (4.1) is sharp for the substitution of some root of unity η into the Burau matrix of β , then by Theorem 6.2 the set I of indices for which Φ_i is pseudo-Anosov, Burau orientable, and satisfies $h_{top}(\Phi_i) = h_{top}(\Phi)$ must be nonempty. Let $K(\beta)$ be the smallest value of k for which there exists $0 < j < k$ with $\lambda = \text{sr}(B(e^{2\pi i j/k}))$. According to Lemma 6.6, $K(\beta)$ is the infimum, over $i \in I$, of twice the power of 2 occurring in the prime factorization of a_i .

Now fix $i \in I$ and let X_i be the supporting surface of Φ_i and $X_{i0} \subset X_i$ a connected component. Adopt the notation of Lemma 6.1 and the definition of a_i . In particular r denotes

the number of punctures and disks deleted from X_{i0} . Because Φ_i is pseudo-Anosov, we must have $r \geq 3$ by the Euler-Poincaré-Hopf formula. Since $a_i \leq \inf_j m_j$ and $\sum_j m_j \leq n/\ell_i$ by definition, we therefore have $a_i \leq n/r\ell_i \leq n/3$. Therefore the power of 2 in a_i is also at most $\frac{1}{3}n$. Since this holds for all $i \in I$, we get $K(\beta) \leq \frac{2}{3}n$, proving Theorem 1.2. ■

Remark 6.11 We remark that if $n = 3n'$ where n' is a power of two, then the braid $\beta_{n'} \in B_n$ constructed after Theorem 6.2 satisfies $K(\beta_{n'}) = \frac{2}{3}n$. Thus for some braids, the bound in Theorem 1.2 is attained.

Acknowledgements: We would like to thank Toby Hall for his plotting software as well as for many helpful conversations, and Jean-Luc Thiffeault for sharing his stimulating unpublished work on the Burau representation and for pointing out that the results in [SKL02] overlap with those in our Theorem 5.1.

REFERENCES

- [AKM65] R. L. Adler, A. G. Konheim, and M. H. McAndrew. Topological entropy. *Trans. Amer. Math. Soc.*, 114:309–319, 1965.
- [Art25] Emil Artin. Theorie der zöpfe. *Abh. Math. Sem. Hamburg*, 4:47–72, 1925.
- [BB05] Joan S. Birman and Tara E. Brendle. Braids: a survey. In *Handbook of knot theory*, pages 19–103. Elsevier B. V., Amsterdam, 2005.
- [Bir74] Joan S. Birman. *Braids, links, and mapping class groups*. Princeton University Press, Princeton, N.J., 1974. Annals of Mathematics Studies, No. 82.
- [Bow71] Rufus Bowen. Entropy for group endomorphisms and homogeneous spaces. *Trans. Amer. Math. Soc.*, 153:401–414, 1971.
- [Bow78] Rufus Bowen. Entropy and the fundamental group. In *The structure of attractors in dynamical systems (Proc. Conf., North Dakota State Univ., Fargo, N.D., 1977)*, volume 668 of *Lecture Notes in Math.*, pages 21–29. Springer, Berlin, 1978.
- [Boy89] Philip Boyland. Notes on dynamics of surface homeomorphisms. *U. of Warwick Preprint Series*, 1989.
- [Boy94] Philip Boyland. Topological methods in surface dynamics. *Topology Appl.*, 58(3):223–298, 1994.
- [Boy99] Philip Boyland. Isotopy stability of dynamics on surfaces. In *Geometry and topology in dynamics (Winston-Salem, NC, 1998/San Antonio, TX, 1999)*, volume 246 of *Contemp. Math.*, pages 17–45. Amer. Math. Soc., Providence, RI, 1999.
- [CLN85] César Camacho and Alcides Lins Neto. *Geometric theory of foliations*. Birkhäuser Boston Inc., Boston, MA, 1985. Translated from the Portuguese by Sue E. Goodman.

- [DGS76] Manfred Denker, Christian Grillenberger, and Karl Sigmund. *Ergodic theory on compact spaces*. Springer-Verlag, Berlin, 1976. Lecture Notes in Mathematics, Vol. 527.
- [Fen83] Roger A. Fenn. *Techniques of geometric topology*, volume 57 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1983.
- [FLP79] Albert Fathi, François Laudenbach, and Valentin Poénaru. *Travaux de Thurston sur les surfaces*, volume 66 of *Astérisque*. Société Mathématique de France, Paris, 1979. Séminaire Orsay, With an English summary.
- [Fra81] John M. Franks. Knots, links and symbolic dynamics. *Ann. of Math. (2)*, 113(3):529–552, 1981.
- [Fri86] David Fried. Entropy and twisted cohomology. *Topology*, 25(4):455–470, 1986.
- [Ful95] William Fulton. *Algebraic topology*, volume 153 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. A first course.
- [GFT06] E. Gouillart, M. D. Finn, and J.-L. Thiffeault. Topological mixing with ghost rods. *Phys. Rev. E*, 73:036311, 2006.
- [KH95] Anatole Katok and Boris Hasselblatt. *Introduction to the modern theory of dynamical systems*, volume 54 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1995. With a supplementary chapter by Katok and Leonardo Mendoza.
- [Kit98] Bruce P. Kitchens. *Symbolic dynamics*. Universitext. Springer-Verlag, Berlin, 1998. One-sided, two-sided and countable state Markov shifts.
- [Kol89] Boris Kolev. Entropie topologique et représentation de Burau. *C. R. Acad. Sci. Paris Sér. I Math.*, 309(13):835–838, 1989.
- [Man74] Anthony Manning. Topological entropy and the first homology group. *Special Year in Dynamical Systems at Warwick University*, 1974.
- [Mil68] John W. Milnor. Infinite cyclic coverings. In *Conference on the Topology of Manifolds (Michigan State Univ., E. Lansing, Mich., 1967)*, pages 115–133. Prindle, Weber & Schmidt, Boston, Mass., 1968.
- [RS75] David Ruelle and Dennis Sullivan. Currents, flows and diffeomorphisms. *Topology*, 14(4):319–327, 1975.
- [Ryk99] E. Rykken. Expanding factors for pseudo-Anosov homeomorphisms. *Michigan Math. J.*, 46(2):281–296, 1999.
- [Sch57] Sol Schwartzman. Asymptotic cycles. *Ann. of Math. (2)*, 66:270–284, 1957.

- [SKL02] Won Taek Song, Ki Hyoungh Ko, and Jérôme E. Los. Entropies of braids. *J. Knot Theory Ramifications*, 11(4):647–666, 2002. Knots 2000 Korea, Vol. 2 (Yongpyong).
- [TF06] J.-L. Thiffeault and M. D. Finn. Topology, braids, and mixing in fluids. *Phil. Trans. R. Soc. Lond.*, 364:3251–3266, December 2006.
- [Thi05] J.-L. Thiffeault. Measuring topological chaos. *Phys. Rev. Lett.*, 94(8):084502, March 2005.
- [Thu88] William P. Thurston. On the geometry and dynamics of diffeomorphisms of surfaces. *Bull. Amer. Math. Soc. (N.S.)*, 19(2):417–431, 1988.

Gavin Band
 Dept. of Mathematics
 University of Liverpool
 Peach St.
 Liverpool L69 7ZL
 U.K.
 g.band@liv.ac.uk

Philip Boyland
 Dept. of Mathematics
 University of Florida
 Little Hall
 Gainesville, FL 32605-8105
 U.S.A.
 boyland@math.ufl.edu